

MAA141Ra21

Monoclonal Antibody to Plasminogen Activator, Urokinase Receptor (uPAR)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: uPAR, RatClonality: Monoclonal

Clone number: H3

Host: Mouse


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 500µg/mL

UOM: 200µg



Sample: Recombinant uPAR, Rat

14

10

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant uPAR (Thr17~Asn212) expressed in *E.coli*.

Accession No.: RPA141Ra01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-TCVP ASQGLRCIQC ESNQDCLVEE CALGQDLCRT TVLREWEDAE ELEVVTRGCA HKEKTNRTMS YRMGSVIVSL TETVCATNLC NRPRPGARGR PFPRGRYLEC ASCTSLDQSC ERGREQSLQC RYPTEHCIEV VTLQSTERSV KDEPYTKGCG SLPGCPGTAG FHSNQTFHFL KCCNFTQCNG GPVLDLQSLP PN



[ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against uPAR. It has been selected for its ability to recognize uPAR in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.