

MAA756Hu21 Monoclonal Antibody to Mucin 5 Subtype AC (MUC5AC) **Organism Species: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

←20kDa

Western Blot

kDa

75

50

37

[PRODUCT INFORMATION]

Immunogen: MUC5AC, Human

Clonality: Monoclonal Clone number: C7

Host: Mouse

Immunoglobulin Type: IgG

Sample: Recombinant MUC5AC, Human

25 20 **Purification:** Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA 15 Concentration: 500µg/mL 10 **UOM:** 200µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant MUC5AC (Tyr4944~His5030) expressed in *E.coli*.

Accession No.: RPA756Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

S-tag and its sequence is listed below.

MHHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF-

YSLEGNT VEHRCQCCQE LRTSLRNVTL HCTDGSSRAF SYTEVEECGC MGRRCPAPGD

TQHSEEAEPE PSQEAESGSW ERGVPVSPMH



[ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against MUC5AC. It has been selected for its ability to recognize MUC5AC in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.