#### MAB867Hu22 Monoclonal Antibody to Epidermal Growth Factor Receptor 2 (EGFR2) Organism Species: Homo sapiens (Human) Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [PRODUCT INFORMATION]

Immunogen: EGFR2, HumanKD2<br/>75Clonality: Monoclonal50Clone number: C131Host: Mouse25Immunoglobulin Type: IgG20Purification: Affinity Chromatography.15Applications: WB, ICC, IHC-P, IHC-F, ELISA10Concentration: 500µg/mL10



Sample: Recombinant EGFR2, Human


### [IMMUNOGEN INFORMATION]

**Immunogen:** Recombinant EGFR2 (Phe376~Gly578) expressed in *E.coli*. **Accession No.:** RPB867Hu01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and GST-tag and its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR

Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003 11271 Richmond Avenue Suite H104. Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Http://www.cloud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F. Wuhan. Hubei 430056, PRC | Toll free: 0086-800-880-0667 | Fax: 0086-27-8425-9551 | Http://www.uscnk.com | E-mail: mail@cloud-clone.us



GSTAIGMKET AAAKFERQHM DSPDLGTLEV LFQGPLGS- FLPES FDGDPASNTA PLQPEQLQVF ETLEEITGYL YISAWPDSLP DLSVFQNLQV IRGRILHNGA YSLTLQGLGI SWLGLRSLRE LGSGLALIHH NTHLCFVHTV PWDQLFRNPH QALLHTANRP EDECVGEGLA CHQLCARGHC WGPGPTQCVN CSQFLRGQEC VEECRVLQGL PREYVNARHC LPCHPECQPQ NGSVTCFG

## [ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against EGFR2. It has been selected for its ability to recognize EGFR2 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

# [<u>CONTENTS</u>]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

# [ <u>STORAGE</u> ]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.