

# MAE361Hu21 Monoclonal Antibody to Na-CI Cotransporter (NCCT)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: NCCT, Human

Clonality: Monoclonal

Clone number: A

Host: Mouse


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 500µg/mL

**UOM**: 200µg



Sample: Recombinant NCCT, Human

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant NCCT (Glu3~Cys146) expressed in *E.coli*.

Accession No.: RPE361Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- ELPTTETP GDATLCSGRF TISTLLSSDE PSPPAAYDSS HPSHLTHSST FCMRTFGYNT IDVVPTYEHY ANSTQPGEPR KVRPTLADLH SFLKQEGRHL

HALAFDSRPS HEMTDGLVEG EAGTSSEKNP EEPVRFGWVK GVMIRC



#### [ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against NCCT. It has been selected for its ability to recognize NCCT in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.