
MAJ323Hu21 Monoclonal Antibody to Cystatin 2 (CST2) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: CST2, Human Clonality: Monoclonal Clone number: G6 Host: Mouse Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 500µg/mL

Sample: Recombinant CST2, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant CST2 (Trp21~Ala141) expressed in E.coli.

Accession No.: RPJ323Hu01

UOM: 200µg

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGS- WSPQEEDRII EGGIYDADLN DERVQRALHF VISEYNKATE DEYYRRLLRV LRAREQIVGG VNYFFDIEVG RTICTKSQPN LDTCAFHEQP ELQKKQLCSF QIYEVPWEDR MSLVNSRCQE A

Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003 11271 Richmond Avenue Suite H104. Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0068 | Http://www.cloud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F. Wuhan, Hubei 430056, PRC | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Http://www.uscnk.com | E-mail: mail@cloud-clone.us

[ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against CST2. It has been selected for its ability to recognize CST2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.