

#### PAA011Po01

Polyclonal Antibody to Brain Derived Neurotrophic Factor (BDNF)
Organism Species: Sus scrofa; Porcine (Pig)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: BDNF, Pig Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100μg



Sample: Recombinant BDNF, Pig

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant BDNF (Pro20~Arg252) expressed in *E.coli*.

Accession No.: RPA011Po01

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSG SEF- P MKEANVRGQG SLAYPGVRTH GTLESVNGPK AGSRGLTSSS SSSLADTFEH VIEELLDEDQ KVRPNEENNK DADMYTSRVM LSSQVPLEPP LLFLLEEYKN YLDAANMSMR VRRHSDPARR GELSVCDSIS EWVTAADKKT AVDMSGGTVT VLEKVPVSKG QLKQYFYETK CNPMGYTKEG CRGIDKRHWN SQCRTTQSYV RAI TMDSKKR IGWRFIRIDT SCVCTI TIKR GR



## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against BDNF. It has been selected for its ability to recognize BDNF in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant BDNF (Pro20~Arg252) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.