PAA098Hu01 Polyclonal Antibody to Matrix Metalloproteinase 10 (MMP10) **Organism Species: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: MMP10, Human **Clonality:** Polyclonal Host: Rabbit Immunoglobulin Type: IgG **Purification:** Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL **UOM:** 100µg Sample: Recombinant MMP10, Human

50 ←45kDa 37 25 20 15 10


[IMMUNOGEN INFORMATION]


Immunogen: Recombinant MMP10 (Phe99~Cys476) expressed in E.coli.

Accession No.: RPA098Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-FS SFPGMPKWRK THLTYRIVNY TPDLPRDAVD SAIEKALKVW EEVTPLTFSR LYEGEADIMI SFAVKEHGDF YSFDGPGHSL AHAYPPGPGL YGDIHFDDDE KWTEDASGTN LFLVAAHELG HSLGLFHSAN TEALMYPLYN SFTELAQFRL SODDVNGIOS LYGPPPASTE EPLVPTKSVP SGSEMPAKCD PALSFDAIST LRGEYLFFKD RYFWRRSHWN PEPEFHLISA FWPSLPSYLD AAYEVNSRDT VFIFKGNEFW AIRGNEVQAG YPRGIHTLGF PPTIRKIDAA VSDKEKKKTY FFAADKYWRF DENSQSMEQG FPRLIADDFP GVEPKVDAVL QAFGFFYFFS GSSQFEFDPN ARMVTHILKS NSWLHC





[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against MMP10. It has been selected for its ability to recognize MMP10 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant MMP10 (Phe99~Cys476) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003 11271 Richmond Avenue Suite H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Http://www.cloud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F, Wuhan, Hubei 430056, PRC | Toll free: 0086-800-880-0667 | Fax: 0066-27-8425-9551 | Http://www.uscnk.com | E-mail: mail@cloud-clone.us