PAA112Hu01 Polyclonal Antibody to Betacellulin (bTC) **Organism Species: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION] Western Blot kDa 75 Immunogen: bTC, Human 50 **Clonality:** Polyclonal Host: Rabbit 37 Immunoglobulin Type: IgG 25 20 **Purification:** Affinity Chromatography. 15 Applications: WB, ICC, IHC-P, IHC-F, ELISA ←11kDa Concentration: 200µg/mL 10 **UOM:** 100µg Sample: Recombinant bTC, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant bTC (Asp32~Tyr111) expressed in E.coli. USCN Accession No.: RPA112Hu01 Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below. MGHHHHHHSGS-DGNSTRSPE TNGLLCGDPE ENCAATTTQS KRKGHFSRCP KQYKHYCIKG RCRFVVAEQT PSCVCDEGYI GARCERVDLF Y

9th Edition (Revised in Jul, 2013)

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against bTC. It has been selected for its ability to recognize bTC in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant bTC (Asp32~Tyr111) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.