

PAA119Po02

Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)

Organism Species: *Sus scrofa*; Porcine (Pig)

*Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PRODUCT INFORMATION ]

**Immunogen:** CD40L, Pig

**Clonality:** Polyclonal

**Host:** Rabbit


**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg



Sample: Recombinant CD40L, Pig

## [ IMMUNOGEN INFORMATION ]

**Immunogen:** Recombinant CD40L (Phe111~Leu261) expressed in *E.coli*.

**Accession No.:** RPA119Po02

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSG SEF- FEMHKGDQDP QIAAHVISEA SSKTASVLQW APKGYTTLST  
NLVTLENGRQ LAVKRQGIYY IYAQVTFCSN RDAAGQAPFI ASLCLRSPSG SERILLRAAN  
THSSKPCGQ QSIHLGGVFE LQPGASVFN VTDPSQVSHG TGFTSFGLLK L

## [ ANTIBODY SPECIFICITY ]

The antibody is a rabbit polyclonal antibody raised against CD40L. It has been

selected for its ability to recognize CD40L in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## **[ CONTENTS ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## **[ QUALITY CONTROL ]**

**Content:** The quality control contains recombinant CD40L (Phe111~Leu261) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.  
5uL per well when used in enhanced chemiluminescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## **[ STORAGE ]**

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.