

#### PAA119Po03

Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)

Organism Species: Sus scrofa; Porcine (Pig)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

Western Blot

#### [ PRODUCT INFORMATION ]

**Immunogen:** CD40L, Pig **Clonality:** Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant CD40L (Phe111~Leu261) expressed in E.coli.

Accession No.: RPA119Po03

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

GST-tag and its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR GSTAIGMKET AAAKFERQHM DSPDLGTLEV LFQGPLGSEF- FEMHKGDQDP QIAAHVISEA SSKTASVLQW APKGYYTLST NLVTLENGRQ LAVKRQGIYY IYAQVTFCSN RDAAGQAPFI ASLCLRSPSG SERILLRAAN THSSSKPCGQ QSIHLGGVFE LQPGASVFVN VTDPSQVSHG TGFTSFGLLK L



# [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against CD40L. It has been selected for its ability to recognize CD40L in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant CD40L (Phe111~Leu261) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.