

PAA144Ra01 Polyclonal Antibody to Vascular Endothelial Growth Factor B (VEGFB) Organism: Rattus norvegicus (Rat) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION]

Immunogen: VEGFB, Rat

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG


Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg

8th Edition (Revised in Jun, 2013)



Sample: Recombinant VEGFB, Rat

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant VEGFB (Ala21~Ser137) expressed in *E.coli*.

USCN Accession No.: RPA144Ra01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-APVSQFDGPS HQKKVVSWID VYARATCQPR EVVVPLSMEL MGNVVKQLVP SCVTVQRCGG CCPDDGLECV PIGOHQVRMQ ILMIQYPSSQ LGEMSLEEHS QCECRPKRKE SAVKPDS



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against VEGFB. It has been selected for its ability to recognize VEGFB in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant VEGFB (Ala21~Ser137) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

Note: The quality control is specifically manufactured as the positive control.

5uL per well when used in enhanced chemilumescent (ECL).

Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.