

PAA221Mu03

Polyclonal Antibody to Transcription factor 20 (TCF20)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [ PRODUCT INFORMATION ]

**Immunogen:** TCF20, Mouse **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

**Immunoglobulin Type**: IgG **UOM**: 100μg

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant TCF20 (Glu1871~Arg1965) expressed in E.coli.

USCN Accession No.: RPA221Mu03

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- ELPLDSNEFW VHEGCILWAN GIYLVCGRLY GLQEALEIAR

EMKCSHCOFA GATI GCYNKG CSFRYHYPCA IDADCI I HEF NESVRCPKHK VRI WR

## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against TCF20. It has been selected for its ability to recognize TCF20 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400


Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [ IMAGES ]



Used in DAB staining on fromalin fixed paraffin- embedded Kidney tissue



Used in Western Blot ,Sample:

Recombinant TCF20,Mouse

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant TCF20 (Glu1871~Arg1965) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

#### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.