PAA280Mu01 Polyclonal Antibody to Calnexin (CNX) Organism Species: Mus musculus (Mouse) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: CNX, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant CNX (Leu263~Pro482) expressed in E.coli.

Accession No.: RPA280Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- LLNDMTPP VNPSREIEDP EDRKPEDWDE RPKIADPDAV KPDDWDEDAP SKIPDEEATK PEGWLDDEPE YIPDPDAEKP EDWDEDMDGE WEAPQIANPK CESAPGCGVW QRPMIDNPNY KGKWKPPMID NPNYQGIWKP RKIPNPDFFE DLEPFKMTPF SAIGLELWSM TSDIFFDNFI ISGDRRVVDD WANDGWGLKK AADGAAEPGV VLQMLEAAEE RP

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against CNX. It has been selected for its ability to recognize CNX in immunohistochemical staining and western blotting.


[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500

Coud-Clone Corp.


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>IMAGES</u>]



Used in DAB staining on fromalin fixed paraffin-embedded Liver tissue

[CONTENTS]



Used in Western Blot, Sample: Recombinant CNX, Mouse

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant CNX (Leu263~Pro482) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.