

#### PAA316Ra01

Polyclonal Antibody to X-Ray Repair Cross Complementing 6 (XRCC6)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

←27kDa

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

Western Blot

kDa 75

50

37

25

20

15

10

Sample: Recombinant XRCC6, Rat

## [ PRODUCT INFORMATION ]

Immunogen: XRCC6, Rat

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100μg



Immunogen: Recombinant XRCC6 (Arg397~Arg603) expressed in E.coli.

Accession No.: RPA316Ra01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- RYTA RKNVSPYFVA LVPQEEELDD QNIQVTPAGF QLVFLPYADD KRKVPFTEKV MANPEQIDKM KAIVQKLRFT YRSDSFENPV LQQHFRNLEA LALDMMESEQ VVDLTLPKVE AIKKRLGSLA DEFKELVYPP GYNPEGKIAK RKADNEGSAS KKPKVELSEE ELKDLFAKGT LGKLTVPALR DICKAYGLKS GPKKQELLEA LSR



### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against XRCC6. It has been selected for its ability to recognize XRCC6 in immunohistochemical staining and western blotting.

# [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant XRCC6 (Arg397~Arg603) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.