

PAA419Ra01**Polyclonal Antibody to Protein Kinase C Gamma (PKCg)****Organism Species: Rattus norvegicus (Rat)****Instruction manual**

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]**Immunogen:** PKCg, Rat**Clonality:** Polyclonal**Host:** Rabbit**Immunoglobulin Type:** IgG**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 200µg/mL**UOM:** 100µg**[IMMUNOGEN INFORMATION]****Immunogen:** Recombinant PKCg (Phe351~Phe614) expressed in *E.coli*.**Accession No.:** RPA419Ra01**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and S-tag and its sequence is listed below.

MHHHHHSSG LVPRGSGMKE TAAKFERQH MDSPDLGTDD DDKAMADIGS
EF-FSFLMVLGKG SFGKVMLAER RGSDELYAIK ILKKDVIVQD DDVDCTLVEK
RVLALGGRGP GGRPHFLTQL HSTFQTPDRL YFVMEYVTGG DLMYHIQQLG KFKEPHAAFY
AAEIAIGLFF LHNQGIHYRD LKLDNVMLDA EGHKITDFG MCKENVFPGS TTRTFCGTPD
YIAPEIIAYQ PYGKSVDWWS FGVLLYEMLA GQPPFDGEDE EELFQAIMEQ TVTYPKSLSR
EAVAICKGFL TKHPGKRLGS GPDGEPTIRA HGFF

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against PKCg. It has been selected for its ability to recognize PKCg in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on formalin fixed
paraffin- embedded brain tissue



Used in Western Blot ,Sample:
Recombinant PKCg,Rat

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant PKCg (Phe351~Phe614) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.