

**PAA440Ra01**

**Polyclonal Antibody to Apoptosis Antagonizing Transcription Factor (AATF)**

**Organism Species: Rattus norvegicus (Rat)**

**Instruction manual**

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## **[ PRODUCT INFORMATION ]**

**Immunogen:** AATF, Rat

**Clonality:** Polyclonal

**Host:** Rabbit


**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg



Sample: Recombinant AATF, Rat

## **[ IMMUNOGEN INFORMATION ]**

**Immunogen:** Recombinant AATF (Ser237~Gln523) expressed in *E.coli*.

**Accession No.:** RPA440Ra01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- SALK NSHKALKALL  
RSLVDLQEEL LFQYPDTRYL VKGTPNAES EEISSEDEL VGEKKKQRKA PPKRKLEMED  
YPSFMAKRSA DFTVYRNRTL QKWHDKTKLA SGKLGKGF GA FERSILTQID HILMDKERLL  
RRTQTKRSAY RVLGKPEPVP EPVAETLPGE PESLPQVPAN AHLKDLDEEI FDDDDFYHQL  
LRELIERKTS SLDPNDQVAM GRQWLAIQKL RSKIRKKVDR KASKGRKLRV HVLSKLLSFM  
APIDHTAMND EARTELYRSL FGQLNRLDAD HGQ

## **[ ANTIBODY SPECIFICITY ]**

The antibody is a rabbit polyclonal antibody raised against AATF. It has been selected for its ability to recognize AATF in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## **[ CONTENTS ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## **[ QUALITY CONTROL ]**

**Content:** The quality control contains recombinant AATF (Ser237~Gln523) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.  
5uL per well when used in enhanced chemiluminescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## **[ STORAGE ]**

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.