

**PAA551Rb51**

**Polyclonal Antibody to Fibroblast Growth Factor 2, Basic (FGF2)**

**Organism Species: *Oryctolagus cuniculus* (Rabbit)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## **[ PRODUCT INFORMATION ]**

**Immunogen:** FGF2, Rabbit

**Clonality:** Polyclonal

**Host:** Cavia


**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg



*Sample: Recombinant FGF2, Rabbit*

## **[ IMMUNOGEN INFORMATION ]**

**Immunogen:** Recombinant FGF2 (Phe26~Ala153) expressed in *E.coli*.

**Accession No.:** RPA551Rb01

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSGSEF-FKDPK RLYCKNGGFF LRIHPDGRVD GVREKSDPHI KLQLQAEERG  
VVSIGVCAN RYLA MKEDGR LLASKCVTDE CFFFERLESN NYNTYRSRKY SSWYVALKRT  
GQYKLGSKTG PGQKAILFLP MSA

## **[ ANTIBODY SPECIFICITY ]**

The antibody is a cavia polyclonal antibody raised against FGF2. It has been selected for its ability to recognize FGF2 in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## **[ CONTENTS ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## **[ QUALITY CONTROL ]**

**Content:** The quality control contains recombinant FGF2 (Phe26~Ala153) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemiluminescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## **[ STORAGE ]**

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.