

PAA585Ga01
Polyclonal Antibody to Cyclin D1 (CCND1)
Organism Species: Chicken (Gallus)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: CCND1, Gallus

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM:** 100µg



Sample: Recombinant CCND1, Gallus

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant CCND1 (Met1~Ile292) expressed in E.coli.

USCN Accession No.: RPA585Ga01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-MEHQLQCCEV ETIRRAYLDA NLLNDRVLQT MLKAEETCSP SVSYFKCVQK EILPYMRKIV ATWMLEVCEE QKCEEEVFPL AMNYLDRFLS FEPLKKSRLQ LLGATCMFVA SKMKETIPLT AEKLCIYTDN SIRPDELLQM ELLLVNKLKW NLAAMTPHDF IEHFLTKMPL AEDTKQIIRK HAQTFVALCA TDVKFISNPP SMIAAGSVVA AVQGLHLGNT NTFLSYQCLT HFLSQVIKCD PDCLRACQEQ IESLLESSLR QAQQHNVSSE TKTVEDEADL SCTPTDVRDV NI



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against CCND1. It has been selected for its ability to recognize CCND1 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant CCND1 (Met1~Ile292) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.