

PAA682Ra01

Polyclonal Antibody to Caspase 12 (CASP12)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: CASP12, Rat

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



Sample: Recombinant CASP12, Rat

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant CASP12 (Met1~Thr244) expressed in E.coli.

Accession No.: RPA682Ra01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- MAAKRTHERD PIYKIKGLAK DMLDGVFDDL MDKNVLNGDE LLKIGEGASL ILSKAENLVE SFFEKTEMAG KIFAGHIANS DKQLSLQFPS DDEEDELQKM FTPSSASESR GKVEDEEMEV NVGVAHASHL MLTVPQGIQS TEVQDSLKLC SRDWFCTMKT ERAEEIYPVM EKEGRTRLAL IICNKKFDYL FDRDDAETDI LNMKELLQNL GYSVVIKENL TAQEMETELM KFAGRPEHQS SDST



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against CASP12. It has been selected for its ability to recognize CASP12 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant CASP12 (Met1~Thr244) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.