



# A90692Mu02 Polyclonal Antibody to Neuropilin 1 (NRP1) Instruction manual

## FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES.

5th Edition (Revised in January, 2013)

### [PRODUCT INFORMATION]

Immunogen: NRP1Clonality: Polyclonal


Host: Rabbit

Species Reactivity: Mouse

Ig type: Rabbit IgG

**Purification:** Antibodies are purified by target protein affinity chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg



Sample: Recombinant mouse NRP1

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant mouse NRP1 (Cys147~Leu265) expressed in E.coli.

Molecular Weight: 14.8kDa

USCN accession No.: P90692Mu02

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- CSQN YTAPTGVIKS PGFPEKYPNS LECTYIIFAP KMSEIILEFE SFDLEQDSNP PGGMFCRYDR LEIWDGFPEV GPHIGRYCGQ KTPGRIRSSS GVLSMVFYTD SAIAKEGFSA NYSVL

11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R. China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com





#### [ ANTIBODY SPEFICITY ]

Anti NRP1 is a rabbit polyclonal antibody raised against mouse NRP1. This antibody has been selected for its ability to recognize mouse NRP1 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

#### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

#### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH 7.4, containing 0.02%Na<sub>3</sub>N, 50% glycerol.

#### [ QUALITY CONTROL ]

Content: The quality control contains recombinant mouse NRP1

(Cys147~Leu265) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not

used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB

0.01%, Na<sub>3</sub>N 0.02%.

#### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.