

PAA786Mu01**Polyclonal Antibody to Ficolin 1 (FCN1)****Organism Species: Mus musculus (Mouse)****Instruction manual**

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]**Immunogen:** FCN1, Mouse**Clonality:** Polyclonal**Host:** Rabbit**Immunoglobulin Type:** IgG**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 200µg/mL**UOM:** 100µg**[IMMUNOGEN INFORMATION]****Immunogen:** Recombinant FCN1 (Arg25~Gly317) expressed in *E.coli*.**Accession No.:** RPA786Mu01**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- RGACPD VKVVGLGAQD
KVVVIQSCPG FPGPPGPKGE PGSPAGRGER GFQGSPGKMG PAGSKGEPGT
MGPPGVKGEK GDTGAAPSLG EKELGDTLCQ RGRPSCKDLL TRGIFLTGWY TIHLPDCRPL
TVLCDMDVDG GGWTVFQRRV DGSIDFFRDW DSYKRGFGNL GTEFWLGN DY
LHLLTANGNQ ELRVDLQDFQ GKGSYAKYSS FQVSEEQEKY KLTLGQFLEG TAGDSLTKHN
NMSFTTHDQD NDANSMNCAA LFHGAWWYHN CHQSNLNGRY LSGSHESYAD GINWGTG

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against FCN1. It has been selected for its ability to recognize FCN1 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on formalin fixed
paraffin-embedded Liver tissue



Used in Western Blot, Sample:
Recombinant FCN1, Mouse

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant FCN1 (Arg25~Gly317) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.