



A90790Mu01

Polyclonal Antibody to Tumor Necrosis Factor Receptor Superfamily, Member 17 (TNFRSF17) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: TNFRSF17 Western Blot **Clonality:** Polyclonal kDa 75 Host: Rabbit 50 **Species Reactivity:** Mouse Ig type: Rabbit IgG 37 **Purification:** Antibodies are purified by 25 target protein affinity chromatography. 20 **Applications:** WB, ICC, IHC-P, IHC-F ← 16kDa 15 Form: Liquid 10 Size: 100µg

Sample: Recombinant mouse TNFRSF17

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant mouse TNFRSF17 (Leu71~Thr184) expressed in

E.coli.

Molecular Weight: 14.0kDa

USCN accession No.: P90790Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-LLRKMNPEAL KDEPQSPGQL DGSAQLDKAD TELTRIRAGD DRIFPRSLEY TVEECTCEDC VKSKPKGDSD HFFPLPAMEE GATILVTTKT GDYGKSSVPT ALQSVMGMFK PTHT

Unique product Superb quality Client favorite Nicest service ($^{\circ}$ ISO9001:2008; ($^{\circ}$ ISO13485:2003; ($^{\circ}$





[ANTIBODY SPEFICITY]

Anti TNFRSF17 is a rabbit polyclonal antibody raised against mouse TNFRSF17. This antibody has been selected for its ability to recognize mouse TNFRSF17 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant mouse TNFRSF17 (Leu71~Thr184) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.