

PAA889Hu01

Polyclonal Antibody to Renin (REN)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: REN, Human **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant REN (Leu67~Glu179) expressed in E.coli.

Accession No.: RPA889Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

 ${\tt MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-LTLG NTTSSVILTN} \\ {\tt YMDTQYYGEI GIGTPPQTFK VVFDTGSSNV WVPSSKCSRL YTACVYHKLF DASDSSSYKH} \\ {\tt YMDTQYYGEI GIGTPPQTFK VVFDTGSSNV WVPSSKCSRL YTACVYHKLF DASDSSSYKH YTACVYHKLF DASDSST DASDSSYKH YTACVYHKLF DASDSST DASDSSYKH YTACVYHKLF DASDSST DASDST DASDSST DASDST DA$

NGTELTLRYS TGTVSGFLSQ DIITVGGITV TQMFGEVTE

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against REN. It has been selected for its ability to recognize REN in immunohistochemical staining and western blotting.

[APPLICATIONS]


Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin-embedded Bowels tissue



Used in Western Blot, Sample: Recombinant REN, Human

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant REN (Leu67~Glu179) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.