

## PAA959Hu01 Polyclonal Antibody to L1-Cell Adhesion Molecule (L1CAM) **Organism Species: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

#### [ PRODUCT INFORMATION ]

Immunogen: L1CAM, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG


**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg





Sample: Recombinant L1CAM, Human

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant L1CAM (Thr218~Gln423) expressed in E.coli.

Accession No.: RPA959Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- TII QKEPIDLRVK ATNSMIDRKP RLLFPTNSSS HLVALQGQPL VLECIAEGFP TPTIKWLRPS GPMPADRVTY QNHNKTLQLL KVGEEDDGEY RCLAENSLGS ARHAYYVTVE AAPYWLHKPQ SHLYGPGETA RLDCQVQGRP QPEVTWRING IPVEELAKDQ KYRIQRGALI LSNVQPSDTM VTQCEARNRH GLLLANAYIY VVQ



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against L1CAM. It has been selected for its ability to recognize L1CAM in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant L1CAM (Thr218~Gln423) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Ruffer: 100mM Tris(nH8 8) 2% SDS 200mM Na(

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol,

BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.