



A90988Mu01 Polyclonal Antibody to Cytochrome P450 2E1 (CYP2E1) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: CYP2E1Clonality: Polyclonal


Host: Rabbit

Species Reactivity: Mouse

Ig type: Rabbit IgG

Purification: Antibodies are purified by target protein affinity chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg



Sample: Recombinant mouse CYP2E1

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant mouse CYP2E1 (Phe378~Ser493) expressed in

E.coli.

Molecular Weight: 14.7kDa

USCN accession No.: P90988Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-FRG YVIPKGTVVI PTLDSLLFDN YEFPDPETFK PEHFLNENGK FKYSDYFKAF SAGKRVCVGE GLARMELFLL LSAILQHFNL KSLVDPKDID LSPVTIGFGS IPREFKLCVI PRS

11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R. China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com





[ANTIBODY SPEFICITY]

Anti CYP2E1 is a rabbit polyclonal antibody raised against mouse CYP2E1. This antibody has been selected for its ability to recognize mouse CYP2E1 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant mouse CYP2E1

(Phe378~Ser493) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not

used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB

0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.