

PAB019Hu01

Polyclonal Antibody to Angiopoietin Like Protein 4 (ANGPTL4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION]

Immunogen: ANGPTL4, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG


Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg

9th Edition (Revised in Jul, 2013)



Sample: Recombinant ANGPTL4, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant ANGPTL4 (His182~Tyr388) expressed in *E.coli*.

Accession No.: RPB019Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and GST-tag and its sequence is listed below.

MSPILGYWKI KGLVQPTRL L LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID
GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV
DFLSKLP EML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK
KRIEAIQID KYLKSSKYIA WPLQG WQATF GGGDHPKSD GSTSGSGHHH HHSAGLVPR
GSTAIGMKET AA AKFERQHM DSPDLGTLEV LFQGPLGSEF- HRLPRDCQE LFQVGERQSG
LFEIQPQGSP PFLVNCKMTS DGGWTVIQRH HDGSVDFNRPWEAYKAGFGD
PHGEFWLGL KVSITGDRN SRLAVQLRDW DGNAELLQFS VHLGGEDTAYSLQTAPVAG
QLGATTVPPS GLSVPFSTWD QDHDLRRDKN CAKSLSGGWW FGTCSHSNLN
GQYFRSIPQQ RQKLKKGIFW KTWGRGRY

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against ANGPTL4. It has been selected for its ability to recognize ANGPTL4 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.