

A91054Mu01
Polyclonal Antibody to Apoptotic Peptidase Activating Factor 1 (APAF1)
Organism: Mus musculus (Mouse)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

[**PRODUCT INFORMATION**]

Immunogen: APAF1, Mouse

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant APAF1, Mouse

[**IMMUNOGEN INFORMATION**]

Immunogen: Recombinant APAF1 (Arg872~Phe1135) expressed in *E. coli*.

USCN Accession No.: P91054Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHS~~G~~SEF- RLKVADCRG HLSWVHGVMF SPDGSSFLTA SDDQ~~T~~IRVWE
TKKVCKNSAI VLKQEIDVVF QENETMVLAV DNIRGLQLIA GKTGQIDYLP EAQVSCCCLS
PHLEYVAFGD EDGAIKIIEL PNNRVFSSGV GHKKAVRHIQ FTADGKTLIS SSED~~S~~VIQVW
NWQTGDYVFL QAHQETVKDF RLLQDSRLLS WSFDGTVK~~V~~W NVITGRIERD FTCHQGT~~V~~LS
CAISSDATKF SSTSADKTAK IWSFDLLSPL HELKGHNGCV RCSAF

Unique product Superb quality Client favorite Nicest service  ISO9001:2008 ;  ISO13485:2003 ; 

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against APAF1. It has been selected for its ability to recognize APAF1 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant APAF1 (Arg872~Phe1135) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.