

PAB206Hu01

Polyclonal Antibody to Mitogen Activated Protein Kinase 14 (MAPK14)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[**PRODUCT INFORMATION**]

Immunogen: MAPK14, Human

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant MAPK14, Human

[**IMMUNOGEN INFORMATION**]

Immunogen: Recombinant MAPK14 (Glu19~Pro279) expressed in *E.coli*.

USCN Accession No.: RPB206Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHH SSSLVPRGSH MASMTGGQQM GRGS-EV PERYQNLSPV GSGAYGSVCA
AFDTKTGLRV AVKKLSRPFQ SIIHAKRTYR ELRLLKHKMH ENVIGLLDVF TPARSLEEFN
DVYLVTHLMG ADLNNIVKCQ KLTDHVVQFL IYQILRGLKY IHSADIIHRD LKPSNLAVNE
DCELKILDFG LARHTDDEMT GYVATRKYRA PEIMLNWMHY NQTVDIWSVG CIMAELLTGR
TLFPGTDHID QLKILRLVVG TPGAELLKKI SSESARNYIQ SLTQMPKMN ANVFIGANP

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against MAPK14. It has been selected for its ability to recognize MAPK14 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN_3 , 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant MAPK14 (Glu19~Pro279) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN_3 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.