



A91367Mu01

Polyclonal Antibody to Vascular Endothelial Growth Factor Receptor 2 (VEGFR2) Organism: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

-33kDa

Western Blot

kDa 75

50

37

25

20

15

10

Sample: Recombinant VEGFR2. Mouse

[PRODUCT INFORMATION]

Immunogen: VEGFR2, Mouse

Clonality: Polyclonal

Host: Rabbit

UOM: 100µg

Immunoglobulin Type: IgG

Purification: Affinity Chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Concentration: 200µg/mL

Solicentiation. 200µg/mil

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant VEGFR2 (Asn46~Thr322) expressed in E.coli.

USCN Accession No.: P91367Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- NTTLQ ITCRGQRDLD WLWPNAQRDS EERVLVTECG GGDSIFCKTL TIPRVVGNDT GAYKCSYRDV DIASTVYVYV RDYRSPFIAS VSDQHGIVYI TENKNKTVVI PCRGSISNLN VSLCARYPEK RFVPDGNRIS WDSEIGFTLP SYMISYAGMV FCEAKINDET YQSIMYIVVV VGYRIYDVIL SPPHEIELSA GEKLVLNCTA RTELNVGLDF TWHSPPSKSH HKKIVNRDVK PFPGTVAKMF LSTLTIESVT KSDQGEYTCV ASSGRMIKRN RT

Unique product Superb quality Client favorite Nicest service 🝳 ISO9001:2008; 🍳 ISO13485:2003 ; $\zeta \in$





[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against VEGFR2. It has been selected for its ability to recognize VEGFR2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant VEGFR2 (Asn46~Thr322) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.