



#### A91499Ra01

## Polyclonal Antibody to Tumor Necrosis Factor Receptor Superfamily, Member 1A (TNFRSF1A) Instruction manual

# FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: TNFRSF1A
Clonality: Polyclonal

Host: Rabbit

Species Reactivity: Rat

Ig type: Rabbit IgG

**Purification:** Antibodies are purified by target protein affinity chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg Western Blot kDa 75 50 37 25 20 ← 21kDa 15

Sample: Recombinant rat TNFRSF1A

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant rat TNFRSF1A (Leu43~Leu194) expressed in E.coli.

Molecular Weight: 18.5kDa

USCN accession No.: P91499Ra01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-LCPQGKYA HPKNNSICCT KCHKGTYLVS DCPSPGQETV CEVCDKGTFT ASQNHVRQCL SCKTCRKEMF QVEISPCKAD MDTVCGCKKN QFQRYLSETH FQCVDCSPCF NGTVTIPCKE KQNTVCNCHA GFFLSGNECT PCSHCKKNQE CMKL

Unique product Superb quality Client favorite Nicest service @ ISO9001:2008; @ ISO13485:2003; CE





#### [ ANTIBODY SPEFICITY ]

Anti TNFRSF1A is a rabbit polyclonal antibody raised against rat TNFRSF1A. This antibody has been selected for its ability to recognize rat TNFRSF1A in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

#### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

#### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH 7.4, containing 0.02%Na<sub>3</sub>N, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant rat TNFRSF1A

(Leu43~Leu194) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading buffer:** 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na<sub>3</sub>N 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.