

PAB563Hu01 **Polyclonal Antibody** To Killer Cell Immunoglobulin Like Receptor 2DS4 (KIR2DS4) Organism Species: Homo sapiens (Human) Instruction manual

9th Edition (Revised in Jul. 2013)

—32kDa

Western Blot

kDa 70

44

33

26

22

18

14

10

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION]

Immunogen: KIR2DS4, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Sample: Recombinant KIR2DS4, Human

Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL **UOM**: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant KIR2DS4 (Gln22~Ser302) expressed in E.coli.

Accession No.: RPB563Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGS- QEGVHRKPS FLALPGHLVK SEETVILQCW SDVMFEHFLL REGKFNNTL HLIGEHHDGV SKANFSIGPM MPVLAGTYRC YGSVPHSPYQ LSAPSDPLDM VIIGLYEKPS LSAQPGPTVQ AGENVTLSCS SRSSYDMYHL SREGEAHERR LPAVRSINGT FQADFPLGPA THGGTYRCFG SFRDAPYEWS NSSDPLLVSV TGNPSNSWPS PTEPSSKTGN PRHLHVLIGT SVVKIPFTIL LFFLLHRWCS DKKNAAVMDQ EPAGNRTVNS EDSDEQDHQE VS



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against KIR2DS4. It has been selected for its ability to recognize KIR2DS4 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant KIR2DS4 (Gln22~Ser302) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control.

Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.