

### PAB586Mu01 Polyclonal Antibody to Integrin Beta 2 (ITGb2) **Organism Species: Mus musculus (Mouse)** Instruction manual

### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



9th Edition (Revised in Jul, 2013)



[IMMUNOGEN INFORMATION]

Immunogen: Recombinant ITGb2 (Ala431~Met652) expressed in E.coli.

#### USCN Accession No.: RPB586Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-ALGFTDTVTV QVRPQCECQC RDQSREQSLC GGKGVMECGI CRCESGYIGK NCECQTQGRS SQELERNCRK DNSSIVCSGL GDCICGQCVC HTSDVPNKEI FGQYCECDNV NCERYNSQVC GGSDRGSCNC GKCSCKPGYE GSACQCQRST TGCLNARLVE CSGRGHCQCN RCICDEGYOP PMCEDCPSCG SHCRDNHTSC AECLKFDKGP FEKNCSVOCA GM

## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against ITGb2. It has been selected for its ability to recognize ITGb2 in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

## [<u>CONTENTS</u>]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant ITGb2 (Ala431~Met652) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

# [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.