

A91812Hu01
Polyclonal Antibody to Beta-Site APP Cleaving Enzyme 2 (bACE2)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: bACE2

Clonality: Polyclonal

Host: Rabbit

Species Reactivity: Human


Ig type: Rabbit IgG

Purification: Antibodies are purified by target protein affinity chromatography.

Applications: WB, ICC, IHC-P, IHC-F

Form: Liquid

Size: 100µg



Sample: Recombinant human bACE2

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant human bACE2 (Met230~Phe396) expressed in *E. coli*.

Molecular Weight: 20.0kDa

USCN accession No.: P91812Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSGS-M QMCGAGLPVA GSGTNGGSLV LGGIEPSLYK GDIWYTPIKE
EWYYQIEILK LEIGGQSLNL DCREYNADKA IVDSGTTLLR LPQKVFDAVV EAVARASLIP

Unique product Superb quality Client favorite Nicest service  ISO9001:2008 ;  ISO13485:2003 ; CE

EFSDGFWTGS QLACWTNSET PWSYFPKISI YLRDENSSRS FRITILPQLY IQPMMGAGLN
YECYRF

[ANTIBODY SPECIFICITY]

Anti bACE2 is a rabbit polyclonal antibody raised against human bACE2. This antibody has been selected for its ability to recognize human bACE2 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02% Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant human bACE2 (Met230~Phe396) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.