

#### PAB817Hu01

Polyclonal Antibody to Versican (VS)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PRODUCT INFORMATION ]

Immunogen: Versican, Human

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



Sample: Recombinant versican, Human

# [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant Versican (Thr2958~Ser3209) expressed in *E.coli*.

Accession No.: RPB817Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGS- TVI TTADEIELEG ATQWPHSTSA SATYGVEAGV VPWLSPQTSE RPTLSSSPEI NPETQAALIR GQDSTIAASE QQVAARILDS NDQATVNPVE FNTEVATPPF SLLETSNETD FLIGINEESV EGTAIYLPGP DRCKMNPCLN GGTCYPTETS YVCTCVPGYS GDQCELDFDE CHSNPCRNGA TCVDGFNTFR CLCLPSYVGA LCEQDTETCD YGWHKFQGQC YKYFAHRRTW DAAERECRLQ GAHLTSILS



## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against versican. It has been selected for its ability to recognize versican in immunohistochemical staining and western blotting.

# [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant versican (Thr2958~Ser3209) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.