PAB973Mu02 Polyclonal Antibody To A Disintegrin And Metalloproteinase With Thrombospondin 1 (ADAMTS1) Organism Species: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: ADAMTS1, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant ADAMTS1 (Thr855~Ser968) expressed in *E.coli*. **USCN Accession No.:** RPB973Mu02

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF -TFSEWV IEEWGECSKT CGSGWQRRVV QCRDINGHPA SECAKEVKPA STRPCADLPC PHWQVGDWSP CSKTCGKGYK KRTLKCVSHD GGVLSNESCD PLKKPKHYID FCTLTQCS

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against ADAMTS1. It has been selected for its ability to recognize ADAMTS1 in immunohistochemical staining and western blotting.

[APPLICATIONS]


Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200

Coud-Clone Corp.

Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>IMAGES</u>]



Western Blot kDa 75 50 37 25 20 15 ← 16kDa 10

Used in DAB staining on fromalin fixed paraffin- embedded Kidney tissue [<u>CONTENTS</u>] Used in Western Blot ,Sample: Recombinant ADAMTS1,Mouse

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant ADAMTS1 (Thr855~Ser968) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.