



A92036Ra02 Polyclonal Antibody to Transferrin (TRF) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: TRF
Clonality: Polyclonal


Host: Rabbit

Species Reactivity: Rat

Ig type: Rabbit IgG

Purification: Antibodies are purified by target protein affinity chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg



Sample: Recombinant rat TRF

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant rat TRF (Ser380~Asn565) expressed in *E.coli*.

Molecular Weight: 25.4kDa

USCN accession No.: P92036Ra02

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

S-tag and its sequence is listed below.

MHHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS- S NGQIECESAE STEDCIDKIV NGEADAMSLD GGHAYIAGQC GLVPVMAENY DISSCTNPQS DVFPKGYYAV AVVKASDSSI NWNNLKGKKS CHTGVDRTAG WNIPMGLLFS RINHCKFDEF FSQGCAPGYK KNSTLCDLCI GPAKCAPNNR EGYNGYTGAF QCLVEKGDVA FVKHQTVLEN





TNGKN

[ANTIBODY SPEFICITY]

Anti TRF is a rabbit polyclonal antibody raised against rat TRF. This antibody has been selected for its ability to recognize rat TRF in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant rat TRF (Ser380~Asn565) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.