

PAC313Mu01

Polyclonal Antibody to Antithrombin (AT)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[**PRODUCT INFORMATION**]

Immunogen: Antithrombin, Mouse

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant Antithrombin, Mouse

[**IMMUNOGEN INFORMATION**]

Immunogen: Recombinant antithrombin (His33~Asn465) expressed in *E. coli*.

Accession No.: RPC313Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSG SEF-HGNPVDD ICIAKPRDIP VNPLCIYRSP GKKATEEDGS EQKVPEATNR
RVWELSKANS RFATNFYQHL ADSKNDNDNI FLSPLSISTA FAMTKLGACN DTLKQLMEVF
KFDTISEKTS DQIHFFFAKL NCRLYRKANK SSDLVSANRL FGDKSLTFNE SYQDVSEVVY
GAKLQPLDFK ENPEQSRVTI NNWVANKTEG RIKDVIPQGA INELTALVLV NTIYFKGLWK
SKFSPENTRK EPFYKVDGQS CPVPMMYQEG KFKYRRAEG TQVLELPFKG DDITMVLILP
KPEKSLAKVE QELPELLQE WLDELSETML VVHMPRFRTD DGFSLKEQLQ DMGLIDLFSF
EKSQPLGIVA GGRDDLYVSD AFHKAFLEVN EEGSEAAAST SVVITGRSLN PNRVTFKANR
PFLVLIREVA LNTIIFMGRV ANPCVN

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against antithrombin. It has been selected for its ability to recognize antithrombin in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN_3 , 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant antithrombin (His33~Asn465) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN_3 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.