

PAC438Hu01

Instruction manual

Polyclonal Antibody to Dymeclin (DYM)

Organism Species: Homo sapiens (Human)

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [ PRODUCT INFORMATION ]

**Immunogen:** DYM, Human **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

**Immunoglobulin Type**: IgG **UOM**: 100μg

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant DYM (Ser546~Asp669) expressed in E.coli.

Accession No.: RPC438Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-SLHHN PNLVYALLYK RDLFEQFRTH PSFQDIMQNI DLVISFFSSR LLQAGAELSV ERVLEIIKQG VVALPKDRLK KFPELKFKYV EEEQPEEFFI PYVWSLVYNS

AVGLYWNPQD IQLFTMDSD

### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against DYM. It has been selected for its ability to recognize DYM in immunohistochemical staining and western blotting.

# [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200




Optimal working dilutions must be determined by end user.

# [ IMAGES ]



Used in DAB staining on fromalin fixed paraffin- embedded Kidney tissue



Used in Western Blot ,Sample: Recombinant DYM,Human

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

# [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant DYM (Ser546~Asp669) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

#### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.