



A92664Hu02 Polyclonal Antibody to Nucleophosmin (NPM) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: NPM Western Blot **Clonality:** Polyclonal kDa Host: Rabbit 75 **Species Reactivity:** Human 50 Ig type: Rabbit IgG 37 **Purification:** Antibodies are purified by 25 target protein affinity chromatography. 20 Applications: WB, ICC, IHC-P, IHC-F 15 Form: Liquid 12kDa **Size:** 100µg 10

Sample: Recombinant human NPM

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant human NPM (Ala204~Trp290) expressed in E.coli.

Molecular Weight: 11.3kDa

USCN accession No.: P92664Hu02

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- AQKSNQN GKDSKPSSTP RSKGQESFKK QEKTPKTPKG PSSVEDIKAK MQASIEKGGS LPKVEAKFIN YVKNCFRMTD QEAIQDLWQW





[ANTIBODY SPEFICITY]

Anti NPM is a rabbit polyclonal antibody raised against human NPM. This antibody has been selected for its ability to recognize human NPM in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant human NPM (Ala204~Trp290) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.