


A93100Hu01
Polyclonal Antibody
To Low Density Lipoprotein Receptor Related Protein Associated Protein 1 (LRPAP1)
Organism: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

[PRODUCT INFORMATION]

Immunogen: LRPAP1, Human
Clonality: Polyclonal
Host: Rabbit
Immunoglobulin Type: IgG
Purification: Affinity Chromatography.
Applications: WB, ICC, IHC-P, IHC-F
Concentration: 200µg/mL
UOM: 100µg



Sample: Recombinant LRPAP1, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant LRPAP1 (Glu90~Leu342) expressed in *E. coli*.
USCN Accession No.: P93100Hu01
Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.
MGHHHHHHSSEF-E LAWKKLKLDG LDEGEKEAR LIRNLNVILA KYGLDGKKDA
RQVTSNSLSG TQEDGLDDPR LEKLWHKAKT SGKFSGEELD KLRWREFLHHK EKVHEYNVLL
ETLSRTEEIH ENVISPSDLS DIKGSVLHSR HTELKEKLRS INQGLDRLRR VSHQGYSTEA
EFEEPRVIDL WDLAQSANLT DKELEAFREE LKHFEAKIEK HNHYQKQLEI AHEKLRHAES
VGDGERVSRS REKHALLEGR TKELGYTVKK HL

Unique product Superb quality Client favorite Nicest service  **ISO9001:2008**;  **ISO13485:2003**; 

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against LRPAP1. It has been selected for its ability to recognize LRPAP1 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant LRPAP1 (Glu90~Leu342) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.