

PAD471Hu01 Polyclonal Antibody to Pappalysin 2 (PAPPA2) Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

← 25kDa

Western Blot

kDal 75

50

37

25

20

15

10

#### [ PRODUCT INFORMATION ]

Immunogen: PAPPA2, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Sample: Recombinant PAPPA2, Human

# Concentration: 200µg/mL **UOM**: 100µg

# [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant PAPPA2 (Gly888~Ala1093) expressed in E.coli.

Accession No.: RPD471Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- GTF RQYVHTASSR RVCDSSGYWT PEEAVGPPDV DQPCEPSLQA WSPEVHLYHM NMTVPCPTEG CSLELLFQHP VQADTLTLWV TSFFMESSQV LFDTEILLEN KESVHLGPLD TFCDIPLTIK LHVDGKVSGV KVYTFDERIE IDAALLTSQP HSPLCSGCRP VRYQVLRDPP FASGLPVVVT HSHRKFTDVE VTPGQMYQYQ VLA



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against PAPPA2. It has been selected for its ability to recognize PAPPA2 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

#### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant PAPPA2 (Gly888~Ala1093) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol,

BPB 0.01%, NaN<sub>3</sub> 0.02%.

# [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.