



A93933Hu02

Polyclonal Antibody to Tumor Necrosis Factor Ligand Superfamily, Member 12 (TNFSF12) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: TNFSF12
Clonality: Polyclonal


Host: Rabbit

Species Reactivity: Human

Ig type: Rabbit IgG

Purification: Antibodies are purified by target protein affinity chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg



Sample: Recombinant human TNFSF12

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant human TNFSF12 (Arg148~Gln247) expressed in

E.coli.

Molecular Weight: 13.5kDa

USCN accession No.: P93933Hu02

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEFELRRQAC- RQI GEFIVTRAGL YYLYCQVHFD EGKAVYLKLD LLVDGVLALR CLEEFSATAA SSLGPQLRLC QVSGLLALRP GSSLRIRTLP WAHLKAAPFL TYFGLFQ

Unique product Superb quality Client favorite Nicest service Q ISO9001:2008; Q ISO13485:2003; CE





[ANTIBODY SPEFICITY]

Anti TNFSF12 is a rabbit polyclonal antibody raised against human TNFSF12. This antibody has been selected for its ability to recognize human TNFSF12 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant human TNFSF12

(Arg148~Gln247) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not

used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB

0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.