



A94348Hu01 Polyclonal Antibody to Solute Carrier Family 3, Member 2 (SLC3A2) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

# [PRODUCT INFORMATION]

| Immunogen: SLC3A2                        | Western Blot |   |         |
|--|--------------|---|---------|
| Clonality: Polyclonal                    | kDa<br>75    |   |         |
| Host: Rabbit                             |              |   |         |
| Species Reactivity: Human                | 50           |   |         |
| <b>Ig type:</b> Rabbit IgG               | 37           |   |         |
| Purification: Antibodies are purified by | 25           |   |         |
| target protein affinity chromatography.  | 20           | _ | ← 18kDa |
| Applications: WB, ICC, IHC-P, IHC-F      | 15           |   | ,       |
| Form: Liquid                             | 10           |   |         |
| <b>Size:</b> 100µg                       | 10           |   |         |

Sample: Recombinant human SLC3A2

### [IMMUNOGEN INFORMATION]

**Immunogen:** Recombinant human SLC3A2 (Leu213~Asp349) expressed in *E.coli*.

Molecular Weight: 16.8kDa

USCN accession No.: P94348Hu01

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-LPAQKWWH TGALYRIGDL QAFQGHGAGN LAGLKGRLDY LSSLKVKGLV LGPIHKNQKD DVAQTDLLQI DPNFGSKEDF DSLLQSAKKK SIRVILDLTP NYRGENSWFS TQVDTVATKV KDALEFWLQA GVDGFQVRD

Unique product Superb quality Client favorite Nicest service 💿 ISO9001:2008; 💿 ISO13485:2003; CE 11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R. China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com





## [ANTIBODY SPEFICITY]

Anti SLC3A2 is a rabbit polyclonal antibody raised against human SLC3A2. This antibody has been selected for its ability to recognize human SLC3A2 in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

## [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH 7.4, containing 0.02%Na<sub>3</sub>N, 50% glycerol.

## [QUALITY CONTROL]

**Content:** The quality control contains recombinant human SLC3A2 (Leu213~Asp349) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading buffer:** 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na<sub>3</sub>N 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Unique product Superb quality Client favorite Nicest service 🙆 ISO9001:2008; 🍳 ISO13485:2003; CE

11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R.China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com