

PAE672Mu01

Polyclonal Antibody to Fascin 2 (FSCN2)

**Organism Species: Mus musculus (Mouse)** 

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

# [ PRODUCT INFORMATION ]

Immunogen: FSCN2, Mouse

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µq



Western Blot

Sample: Recombinant FSCN2, Mouse

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant FSCN2 (Gln8~Asn278) expressed in E.coli.

Accession No.: RPE672Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF-QVL KIQFGLVNDA DRYLTAESFG FKVNASAASL KRKQIWVLEP DPGQGTAVLF RSSHLGRYLS AEEDGRVACE MDQPGRDCRF LVLPQPDGRW VLQSEPHGRF FGGIEDRLSC FATAISPAEL WTVHLAIHPQ AHLLSVSRRR YVHLCLQEDE MAADGDMPWG VDALVTLIFQ SRRYCLKSYD SRYLRSDGRL VWEPEAHACY TLEFKAGKLA FKDCDGRYLA PVGPAGTLKA GRNTRPSKDE LFDLEQSHPQ VVLVAANRRY ISVRQGIN



## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against FSCN2. It has been selected for its ability to recognize FSCN2 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

#### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant FSCN2 (Gln8~Asn278) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100 mM Tris(pH8.8), 2% SDS, 200 mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.