

PAE677Mu01

Polyclonal Antibody to FKBP7 Binding Protein 7 (FKBP7)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: FKBP7, Mouse

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant FKBP7 (Gln20~His213) expressed in *E.coli*.

Accession No.: RPE677Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSG SEF-Q GQTKEESTEE VKIEVLHRPE NCSKTSRKGD LLNAHYDGYL
AKDGSKFYCS RTQDEGHPKW FVLGVGHVIK GLDIAMDMC PGEKRKVIIP PSFAYGKEGY
AEGKIPP NAT LMFEIELYAV TKGPRSIETF KQIDTDNDRQ LSKAEIELYL QKDFEKDANP
RDKSYQKAVL EDIFKKNDHN GDGFISPKEY NVH

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against FKBP7. It has been selected for its ability to recognize FKBP7 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[**IMAGES**]



Used in DAB staining on formalin fixed
paraffin- embedded lung tissue



Used in Western Blot ,Sample:
Recombinant FKBP7,Mouse

[**CONTENTS**]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[**QUALITY CONTROL**]

Content: The quality control contains recombinant FKBP7 (Gln20~His213) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[**STORAGE**]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.