



# A98123Hu02 Antibody to Fumarylacetoacetate Hydrolase (FAH) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

←27kDa

Western Blot

kDa

75 50

37

25

20

15

#### [ PRODUCT INFORMATION ]

Immunogen: FAH
Clonality: Polyclonal

Host: Rabbit

Organism: Human

**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

Concentration: 200µg/mL

**UOM:** 100µg



### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant human FAH (Val189~Ser419) expressed in *E.coli*.

**USCN Accession No.:** P98123Hu02

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF- VY GACKLLDMEL EMAFFVGPGN RLGEPIPISK AHEHIFGMVL MNDWSARDIQ KWEYVPLGPF LGKSFGTTVS PWVVPMDALM PFAVPNPKQD PRPLPYLCHD EPYTFDINLS VNLKGEGMSQ AATICKSNFK YMYWTMLQQL THHSVNGCNL RPGDLLASGT ISGPEPENFG SMLELSWKGT KPIDLGNGQT RKFLLDGDEV IITGYCQGDG YRIGFGQCAG KVLPALLPS





### [ANTIBODY SPECIFITY]

Anti FAH is a rabbit polyclonal antibody raised against human FAH. This antibody has been selected for its ability to recognize human FAH in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% Na<sub>3</sub>N, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant human FAH(Val189~Ser419) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na<sub>3</sub>N 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.