

PAJ279Mu01**Polyclonal Antibody****To Dolichyl Diphosphooligosaccharide Protein Glycosyltransferase (DDOST)****Organism Species: Mus musculus (Mouse)*****Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]**Immunogen:** DDOST, Mouse**Clonality:** Polyclonal**Host:** Rabbit**Immunoglobulin Type:** IgG**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 200µg/mL**UOM:** 100µg**[IMMUNOGEN INFORMATION]****Immunogen:** Recombinant DDOST (Gly29~Pro412) expressed in *E.coli*.**Accession No.:** RPJ279Mu01**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHS G SEF-GP RTLVL DNLN VRDTHSLFFR SLKDRGFELT FKTADDP SLS
LIKYGFLYD NLIIFSPSVE DFGGNIN VET ISAFIDGGGS VLVAASSDIG DPLREL GSEC
GIEFDEEKTA VIDHHNYDVS DLGQHTLIVA DTENLLKAPT IVGKSSLNPI LFRGVGMVAD
PDNPLVLDIL TGSSTSYSFF PDKPITQYPH AVGRNTLLIA GLQARNNARV IFSGSLDFFS
DAFFNSAVQK ATPGAQRYSQ TGN YELAV AL SRWVFKEEGV LRVGPVSHHR
VGEMAPPNAY TVTDLVEYSI IIEQLSNGKW VPFDGDDIQL EFVRIDPFVR TFLKRKGGKY
SVQFKLPDVY GVFQFKVDYN RLG YTHLYSS TQVSVRPLQH TQYERFIPSA YP

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against DDOST. It has been selected for its ability to recognize DDOST in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500


Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on formalin fixed
paraffin- embedded liver tissue



Used in Western Blot ,Sample:
Recombinant DDOST,Mouse

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant DDOST (Gly29~Pro412) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.