

PAJ829Hu03
Polyclonal Antibody
To Phosphoinositide-3-Kinase Catalytic Beta Polypeptide (PIK3Cb)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

-23kDa

Western Blot

75

50

37

25

20

15

10

### [ PRODUCT INFORMATION ]

Immunogen: PIK3Cb, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg Sample: Recombinant PIK3Cb, Human

# [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant PIK3Cb (Arg321~Pro507) expressed in E.coli.

Accession No.: RPJ829Hu03

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSG SEF- RIISHVWENN NPFQIVLVKG NKLNTEETVK VHVRAGLFHG TELLCKTIVS SEVSGKNDHI WNEPLEFDIN ICDLPRMARL CFAVYAVLDK VKTKKSTKTI NPSKYQTIRK AGKVHYPVAW VNTMVFDFKG QLRTGDIILH SWSSFPDELE EMLNPMGTVQ TNPYTENATA LHVKFPENKK QPYYYPP



### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against PIK3Cb. It has been selected for its ability to recognize PIK3Cb in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant PIK3Cb (Arg321~Pro507) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.