

PAK477Hu01 Polyclonal Antibody to Transmembrane Protein 27 (TMEM27) **Organism Species: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

—16kDa

Western Blot

kDa 75

50

37

25

20

15

10

[PRODUCT INFORMATION]

Immunogen: TMEM27, Human

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Sample: Recombinant TMEM27, Human

Concentration: 200µg/mL **UOM**: 100µg [IMMUNOGEN INFORMATION]

Immunogen: Recombinant TMEM27 (Cys17~Asp137) expressed in E.coli.

Accession No.: RPK477Hu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- CQPG AENAFKVRLS IRTALGDKAY AWDTNEEYLF KAMVAFSMRK VPNREATEIS HVLLCNVTQR VSFWFVVTDP SKNHTLPAVE VQSAIRMNKN RINNAFFLND **QTLEFLKIPS TLAPPMD**



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against TMEM27. It has been selected for its ability to recognize TMEM27 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant TMEM27 (Cys17~Asp137) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control.

Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.