# PAL057Mu01 Polyclonal Antibody to PTEN Induced Putative Kinase 1 (PINK1)

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES


[ PRODUCT INFORMATION ] Western Blot kDa 75 Immunogen: PINK1, Mouse 50 **Clonality:** Polyclonal 37 Host: Rabbit 25 Immunoglobulin Type: IgG 20 **Purification:** Affinity Chromatography. 15 Applications: WB, ICC, IHC-P, IHC-F, ELISA 10 Concentration: 200µg/mL **UOM:** 100µg Sample: Recombinant PINK1, Mouse

#### [IMMUNOGEN INFORMATION]

Immunogen: Recombinant PINK1 (Gly310~Leu566) expressed in E.coli. USCN Accession No.: RPL057Mu01 Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below. MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-G RTLFLVMKNY PCTLRQYLEE QTPSSRLATM MTLQLLEGVD HLVQQGIAHR DLKSDNILVE WDSDGCPWLV ISDFGCCLAD QHVGLRLPFN SSSVERGGNG SLMAPEVSTA HSGPSAVIDY SKADTWAVGA IAYEIFGLAN PFYGQGSAHL ESRSYQEAQL PEMPESVPPE ARRLVRSLLQ REASKRPSAR LAANVLHLSL WGEHLLALKN LKLDKMIAWL LQQSAATLLA DRLREKSCVE TKLQMLFLAN LECEAL

9th Edition (Revised in Jul, 2013)

## **Organism Species: Mus musculus (Mouse)** Instruction manual



### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against PINK1. It has been selected for its ability to recognize PINK1 in immunohistochemical staining and western blotting.

#### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

### [<u>CONTENTS</u>]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant PINK1 (Gly310~Leu566) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

### [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.